



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

That the above formula does not give the volume bounded by $f(x, y, z) = 0$, the xy -plane and a cylinder whose elements are parallel to the z -axis, may be readily seen by applying it to the plane $z = c$, in which case it gives a result one-third as large as the correct result.

Also solved by the Proposer.

239 (Number Theory) [March, 1916]. Proposed by HAROLD T. DAVIS, Colorado Springs, Colorado.

Give a general method for determining the solution in integers of the equation

$$x^r - 10xy - (n + 1) + y = 0,$$

where n and r are positive integers.

SOLUTION BY ELIJAH SWIFT, University of Vermont.

Solving for y , $y = \frac{x^r - (n + 1)}{10x - 1}$, which must be an integer. Since the denominator is prime to 10 and the numerator integral, y will be an integer if $\frac{10^r x^r - (n + 1) \cdot 10^r}{10x - 1}$ is. Dividing algebraically the remainder is $1 - (n + 1) \cdot 10^r$. If then $\frac{10^r(n + 1) - 1}{10x - 1}$ is an integer, so is y . Hence the general process (perhaps not that desired) is the following: form $10^r(n + 1) - 1$ and factor it. Equate $10x - 1$ to any factor whose last digit is 9, and we have an integral solution.

261 (Number Theory) [March, 1917]. Proposed by NORMAN ANNING, Chilliwick, B. C.

Show that for any positive integer n (excluding powers of 2) positive integers $a_1, a_2, a_3, \dots, a_k$ which are less than $n/2$ can be chosen in such a way that

$$2^k \cos(a_1\pi/n) \cos(a_2\pi/n) \cos(a_3\pi/n) \cdots \cos(a_k\pi/n) = 1.$$

SOLUTION BY C. F. GUMMER, Queen's University.

Since n is not a power of 2, it is of the form $(2k + 1)l$. The equation

$$\cos(2k + 1)x - \cos(2k + 1)\alpha = 0$$

has $2k + 1$ distinct roots in $\cos x$, when $\cos(2k + 1)\alpha \neq 1$, the roots being

$$\cos\left(\alpha + \frac{2i\pi}{2k + 1}\right), \quad i = 0, 1, \dots, 2k.$$

Also $\cos(2k + 1)x = 2^{2k} \cos^{2k+1} x - \dots$, the absolute term being zero. Hence,

$$2^{2k} \prod_{i=0}^{2k} \cos\left(\alpha + \frac{2i\pi}{2k + 1}\right) = \cos(2k + 1)\alpha.$$

By taking the limit of each side when $\alpha \rightarrow 0$,

$$2^{2k} \prod_{i=0}^{2k} \cos \frac{2i\pi}{2k + 1} = 1,$$

that is,

$$\left\{ 2^k \prod_{i=0}^k \cos \frac{2i\pi}{2k + 1} \right\}^2 = 1,$$

or

$$2^k \prod_{i=0}^k \cos \frac{2i\pi}{2k + 1} = \pm 1.$$

By taking $j = 2i$, when $i \leq k/2$ and $j = 2k + 1 - 2i$ when $i > k/2$, we get

$$2^k \prod_{j=1}^k \cos \frac{j\pi}{2k + 1} = +1,$$

which takes the required form if $a_j = jl$.